



Procidic2[®]

Broad Spectrum Bactericide and Fungicide Compound

Your solution for powdery mildew, bud rot and root rot

Procidic2 concentrate is a fungicide and bactericide labeled specifically for use on cannabis, marijuana and hemp. Procidic2 has been approved under Washington State Department of Agriculture Organic Food Program for use in organic production.

Working systemically, Procidic2 is rapidly absorbed into the plant and moves toward the apex allowing new tissues to be free of infection and reducing the risk of future growth of bacteria and fungi. Apply Procidic2 in regularly scheduled spray programs or in tank mixes.

▶ See back for more details or visit us at www.sarasotagg.com

Prevents and Controls

- **Powdery Mildew**
- **Bud Rot**
- **Root Rot**
- Fungicide and bactericide dual mode of action
- Best used as a preventative or at first sign of disease
- For use in organic production
- Use on plants grown indoors, in greenhouses or on field plants
- Works on contact
- Works systemically
- No re-entry waiting period required
- Use up to day of harvest



Greenspire Global, Inc. represents that this product is a minimum risk pest control product and qualifies for exemption from EPA registration under FIFRA 25 (b).

Procidic²®

Broad Spectrum Bactericide and Fungicide Compound

Recommended Usage Rates & Instructions

1. Procidic2 is best used as a preventative, although repeat curative applications can be very successful before week 6 of flower
2. Use reverse osmosis water or water under 200ppm
3. Adjust water pH to 5.5-6.0 BEFORE adding Procidic2
4. Apply Procidic2 in a regularly scheduled spray program
5. Use up day of harvest
6. No re-entry waiting period required
7. Use a higher dosage of Procidic2 during periods of high humidity
8. Do not mix with salt-based or high-alkali reaction products



Available in:
32 fl oz, 1 gallon and 2 ½ gallon sizes

Use Rates

Crop	Target Disease	Use Rate	Recommendations
Greenhouse, Indoor-grown or Field-grown Cannabis	Powdery Mildew (<i>Sphaerotheca macularis</i>)	Preventative: 0.5-3.0 fl oz/gallon (15-90 ml/gallon)	Spray when favorable weather conditions are expected, or when disease inoculum is present. Spray every 5 to 10 days.
	Gray Mold/Bud Rot (<i>Botrytis cinerea</i>)	Curative: 1-3.5 fl oz/gallon (25-100 ml/gallon)	Upon discovery of disease, spray every 3 to 5 days. Repeat until control is attained.
Greenhouse, Indoor-grown or Field-grown Cannabis	Root Rot (<i>Pythium sp</i> , <i>Fusarium sp</i> , <i>Phytophthora sp</i> , <i>Rhizoctonia sp</i>)	Preventative: 0.4-0.5 fl oz/gallon (12-15 ml/gallon)	Root drench preventatively every 7 to 14 days.
		Curative: 0.5-1 fl oz/gallon (15-25 ml/gallon)	Upon discovery of disease, root drench every 5 days. Repeat until control is attained.

► For more information go to www.sarasotagg.com or call (800) 550-6259

Procidic² is distributed through a licensing agreement with:
Sarasota Green Group • 5550 Wild Rose Lane, Suite 400 West Des Moines, IA 50266 • 800-550-6259

Procidic² is a registered trademark product manufactured by GreenSpire Global, Inc.